

SCLERAL GRAFTS IN OPHTHALMIC SURGERY. A REVIEW

Dítě Jakub, Netuková Magdaléna, Procházková Alexandra,
Poláchová Martina, Krivosheev Katarína, Studený Pavel

Department of Ophthalmology, 3rd Faculty of Medicine, Charles
University and Královské Vinohrady University Hospital in Prague,
Czech Republic

The authors of the study declare that no conflict of interests exists in the compilation, theme and subsequent publication of this professional communication, and that it is not supported by any pharmaceuticals company. The study has not been submitted to any other journal or printed elsewhere.

Submitted to the editorial board: August 29, 2023

Accepted for publication: December 20, 2023

Available on-line: February 14, 2024



MUDr. Jakub Dítě
Oftalmologická klinika 3. lékařské
fakulty Univerzity Karlovy a Fakultní
nemocnice Královské Vinohrady
Šrobárova 1150/50
Praha 10
E-mail: jakub.dite@gmail.com

SUMMARY

Aim: To summarize the history and current trends in the use of scleral grafts in ophthalmology.

Materials and methods: We conducted a review of the literature through the MEDLINE and Cochrane Library databases. The search terms were "sclera", "graft", and "surgery". The search resulted in 1596 articles, of which we evaluated 192 as relevant. The relevant articles were sorted chronologically and according to the method of using scleral grafts, which enabled the development of a review article.

Results: The sclera has been routinely used in ophthalmology since the 1950s in many different indications. Some of these indications have become practically obsolete over time (for example, use in the surgical management of retinal detachment), but a large number still find application today (especially use in glaucoma or oculoplastic surgery, or as a patch for a defect in the sclera or cornea).

Conclusion: Even though allogeneic sclera is currently used less frequently in ophthalmology compared to other tissue banking products and the range of its indications has partially narrowed, it remains a useful material due to its availability and properties.

Key words: sclera, graft, surgery, ophthalmology, review

Čes. a slov. Oftal., 80, 2024, No. 5, p. 231–238

INTRODUCTION

The sclera is formed during embryonic development from the cells of the neural crest, and together with the cornea it forms the tunica fibrosa oculi. It constitutes approximately five sixths of the surface of the eyeball, and similarly to the cornea is composed especially of type 1 collagen fibers, which, however, in the case of the sclera are not regularly organized, with the result that unlike the cornea it is not transparent. It also contains only a relatively small quantity of cells and blood vessels. It is made up of four layers – the episclera, the stroma, the lamina fusca and the endothelium. The thickness of the sclera is 0.3–1.0 mm, reaching the highest values in the region of the posterior pole, and the lowest in the surrounding area of the tendons of the oculomotor muscles.

With regard to its structure, which gives it strength as well as relatively low antigenicity and low susceptibility to rejection thanks to the small quantity of blood vessels and cells, the sclera has long been considered promising material. It can be used in ocular surgery, not only in autologous but also in allogeneic transplantation.

MATERIAL AND METHODS

During the compilation of this review article we conducted an analysis of the literature through the MEDLINE and Cochrane Library databases. The search terms were "sclera", "graft", and "surgery". The search resulted in 1596 articles, of which we evaluated 192 as relevant. The relevant articles were sorted chronologically and according to the method of using scleral grafts, which enabled the compilation of this review article. The sources of the articles we found were also investigated for the purpose of potentially finding further relevant sources of literature which were not identified during the original search.

THE FIRST MENTIONS OF THE USE OF THE SCLERA

The first mention of the use of the sclera in ophthalmic surgery originates from the 1940s, when Costa published an article on the subject of subconjunctival scleral grafts [1], and Larsson described the treatment of corneal per-

foration with the aid of a patch from an autologous sclera [2]. In the following years, further authors including Kurz [3,4] focused on compensating for a defect in the cornea by means of a scleral graft. Another possible use of the sclera was described at the beginning of the 1950s by Paufigue, who made use of a scleral patch in the surgical treatment of scleromalacia [5].

Originally, autologous [2] or fresh allogeneic tissue [6,7] was used exclusively. However, the 1950s and 60s saw a period of precipitous development of tissue banking, which thanks to new methods of conservation and storage of tissues enabled a previously unimaginable expansion of the use of allogeneic tissues in ophthalmic surgery, which related also to the sclera. In 1961 Payrau was the first to publish the use of a lyophilized sclera [8]. In 1962 Rodriguez-Vasquez then described the use of a sclera conserved in xylol [9], in 1964 Wilson published the use of a sclera conserved in glycerol [10], and in 1965 Knobloch described the use of a sclera conserved in ethanol [11]. These new procedures enabled a further dramatic expansion of the possibilities of using the sclera in ophthalmic surgery.

CORNEAL DEFECTS

As mentioned in the introduction, one of the oldest available publications describing the use of the sclera in ophthalmology is Larsson's article from 1948, which outlines its successful use for the closure of a corneal defect upon the background of a corneal ulcer [2]. In the 1950s his study was followed up by further authors including Kurz [3,4]. Even despite the fact that this is a rather rare method it still has its application, and in very complicated cases or in case of a lack of other options it is still used to this day for the closure of corneal perforations of various etiology [12–22]. An interesting fact is that after implantation into the cornea, the sclera has a tendency towards translucency [3,13,15]. In addition, in contrast with corneal grafts, only a thin scleral patch is required for closing the defect, thanks to which, according to Prydal, more rapid visual rehabilitation takes place upon use on the corneal periphery, at the same time with lesser induced postoperative astigmatism [15].

DEFECTS AND THINNING OF THE SCLERA, INCLUDING SCLEROMALACIA

The surgical treatment of scleromalacia with the aid of a scleral patch was described by Paufigue as early as in 1953 [5], and his work was progressively followed up also by other authors [7,23]. Over the course of the following years, methods of treating immune-conditioned [24–30], surgically induced – most often as a consequence of a pterygium removal surgery [8,28,31–40], infectious [41–46], post-traumatic [40,47] and post-irradiation [35,37,48–50] scleral thinning were successfully described.

In 1959 Kanagasundaram published the possibility of using a scleral graft in the treatment of a traumatic rupture of the eyeball [6]. In this indication also, the sclera

continued to be used frequently [51,52]. Karaca described good results in the treatment of penetrating ocular trauma with the use of only an autologous lamellar scleral graft [22]. Turaga again described the successful use of a scleral patch in a patient with Marfan syndrome who was suffering from recurrent spontaneous ruptures of the eyeball [53].

Reconstruction of the orbital wall following extensive resection procedures, most often due to melanoma of the uvea, became another common indication for the use of scleral or sclerocorneal grafts. The first to describe this technique was Tarkkanen in 1967, in an surgery of melanoma on the limbus [54], after which other authors also described the possibility of reconstruction, for example following corneoscleroidectomy for melanoma of the ciliary body [55,56] or following sclerochoroidectomy for melanoma of the choroid [57]. It is possible to use an analogous procedure to treat squamous cell carcinoma [58] or benign tumors – for example limbal dermoid [59,60]. Rare indications to have been described include reconstruction of the eye following resection of Kaposi's sarcoma of the anterior segment of the eye in a child following bone marrow transplant for acute lymphoblastic leukemia [61].

At present the use of a scleral graft for treating a defect or thinning of the sclera remains a highly relevant technique. Alternatively, it is possible to use the fascia lata femoris [27,62], periosteum [45], perichondrium [63,64], pericardium [65–67], temporal fascia [68], buccal mucosa [49,69], tarsoconjunctival flap [70], amnion [71,72] or cornea [71,73–75].

SCLEROPLASTY

The option of treating progressive myopia in children with the aid of scleroplasty using a scleral graft was first published by Borley in 1958, and his work was then followed up by further authors [76–80]. In 2018 Xue published good results upon the use of an allogeneic sclera treated with the crosslinking method [81].

Other biological materials are also used as an alternative in scleroplasty, for example preparations from porcine (pig) skin [82], the dura mater [83] or lyophilized fascia lata femoris [84], as well as synthetic materials (Gore-Tex) [80].

VITREORETINAL SURGERY

Based on the study by Rodriguez-Vasquez from 1962 [9], the sclera was used for a long time as an alternative material in retinal detachment surgery with an external approach, and in the following years it was successfully used in this indication with various modifications also by further authors [10,11,85–89].

With regard to the fact that today the gold standard in the treatment of retinal detachment is considered to be pars plana vitrectomy [90], and in the case of surgery with an external approach synthetic materials are used

in the overwhelming majority of cases – including silicone, hydrogel, and Gore-Tex [91,92], the use of the sclera in this indication is now practically obsolete.

Another innovative use of the sclera in vitreoretinal surgery was published by Shah in 2017. This describes the possibility of treating macular edema caused by an optic nerve pit through the implantation of a scleral graft for the purpose of closing the pit [93].

OCULOPLASTIC SURGERY

Within the field of oculoplastic surgery the sclera found its first use in 1968, when Helveston published the results of the treatment of extrusion of an orbital implant in patients following enucleation by means of covering with a scleral patch [94]. A large number of studies have been published on this subject over the course of the years [95–103], and the sclera is still used to this day in this indication in various modifications. With regard to the highly positive results of these operations and relatively frequent extrusions, mainly in the case of certain types of orbital implants (especially implants made of hydroxyapatite), Soll published a study in 1974 in which he described a very good outcome upon covering orbital implants with a scleral patch already in the primary operation [104]. This technique also became established, and in the following years several other authors also published their results [105–110]. A less frequent possibility of the use of the sclera in patients following enucleation was described in 1983 by Smith, who used it as a filling material upon atrophy of the orbital tissues [111].

Alternative materials used for covering implants are the fascia lata femoris [101], bovine pericardium [107,112], labial mucosa [113] or Vicryl mesh [114].

Another possibility for use of the sclera in oculoplastic surgery was described by Bodian in 1968 [115]. In frontalis suspension surgery for the purpose of correcting ptosis he used a fresh or conserved sclera as an alternative to the fascia lata femoris. His work was followed up in 1975 by Helveston, who recommended reinforcing the sclera with a non-absorbable suture [116].

However, at the present time substantially more commonly used materials in frontalis suspension surgery include especially the aforementioned autogenous or allogeneic fascia lata femoris [117–119] and synthetic materials such as silicone, Gore-Tex or polypropylene [119–121].

Another indication for use of the sclera is replacement of the posterior lamella of the eyelid upon treatment of cicatricial entropion, which was first published by Tenzel and Rubenzik in 1975 [122,123], and followed on from in the following years also by several other authors [124–126]. It is possible to use an analogous procedure, namely replacement of the posterior lamella of the eyelid with a sclera, also for retraction of eyelids of various etiology (including thyroid eye disease), which was first published by Crawford in 1976 [127], and subsequently

by several more authors [124,128–136], or more extensive defects following resection procedures, for example as part of modification of the Cutler-Beard procedure [137,138], or following resection due to severe ligneous keratoconjunctivitis [139].

In general the consensus predominates among authors that in the case of replacement of part of the eyelid with a sclera it is necessary to use an implant of larger dimensions than the specific defect, because contraction of the scleral graft will take place over the course of time [123,129].

Use of the sclera upon treating retraction of the eyelids upon a background of thyroid eye disease in particular continues to be a highly topical and widely used method. Alternative materials that can be used successfully to replace the posterior lamella of the eyelid include also synthetic materials such as porous polyethylene (Medpor) [136], free tarsoconjunctival flap [135], mucosa of the hard palate [135,136], tragal cartilage (taken from the ear tragus, usually as an autotransplant) [136,138] or acellular dermal matrix (AloDerm) [136]. In patients following resection of the eyelids, an autologous graft from the periosteum of the orbital rim is frequently used, which at the same time provides fixation to the bone and is thereby advantageous for reconstruction especially of the lateral canthus [140].

GLAUCOMA SURGERY

Nesterov was the first to use the sclera in the surgical treatment of glaucoma. In his study from 1978 he described the method of implantation of scleral strips into the supraciliary space, if necessary in combination with the performance of cyclodialysis [141].

In 1987 Fredman published the possibility of preventing extrusion of a Molteno implant by means of covering with a sclera [142], which is a commonly used method to this day in the covering of glaucoma implants, published in a number of different variations also by other authors [143–154]. Lam described the covering of an Ahmed valve in an analogous manner [143], Aslanides proposed the use of an autologous sclera in this indication due to better availability [144], and Zeppa published a modification with fixation of the sclera with the aid of tissue adhesive [155].

Other materials to have been used successfully in the covering of glaucoma implants are the pericardium [145,148,150,156–159], dura mater [145,146], cornea [152,154,156,157,160,161], tragal perichondrium [162], collagen implant Ologen [163], fascia lata femoris [157] or amniotic membrane [157,158]. If necessary it is possible to insert the implant into a scleral tunnel, in which case no further covering implant is required [151,157,164,165]. In recent times the cornea especially has demonstrated itself to be an advantageous alternative in this indication due to its good availability (as a waste product of refractive surgery) and its good cosmetic effect [152].

Another very common indication for use of the sclera in glaucoma surgery is the treatment of postoperative complications. In 1991 Melamed published the method of treating persistent leakage from a filtration bleb with the aid of a scleral patch [166], which is a technique used to this day in the case of leakage or excessive filtration, and has been described by several other authors [167–179]. In 2009 King proposed a modified technique with only a small scleral patch fixed with tissue adhesive [176]. A scleral patch can be used effectively also to treat extrusion of a glaucoma implant [180,181], and if necessary it can be used in revision for postoperative blebitis with endophthalmitis [182].

In the treatment of complications with a filtration bleb, alternatives to the sclera used successfully in the past also include the fascia lata femoris [183], cornea [183,184], dura mater [183], pericardium [183], acellular dermal graft [183], buccal mucosa [183], tragal perichondrium [183], amniotic membrane [183], polytetrafluorethylene [183] or Ologen [183].

Another proposed option for use of the sclera in glaucoma surgery is its implantation in deep sclerectomy, which was described in the past by Devloo and subsequently by Mousa [185,186].

OTHER INDICATIONS

Scleral or sclerocorneal grafts can also be used in the treatment of postoperative or post-traumatic epithelial ingrowth into the anterior chamber of the eye, which was first published by Naumann in 1992 [187] and subsequently by Forster and Rummelt, who also used this technique successfully [188,189].

A relatively new method was described in strabismus surgery, which was published by Thorisdottir in 2018. In complicated cases a scleral strip is used successfully for configuring the length of the oculomotor muscle [190,191].

STATISTICAL DATA

In order to demonstrate the frequency of use of the sclera in various indications in real conditions of modern ophthalmic surgery, here we present data from Catalonia, which was published in a review article from the year 2020 by Sabater-Cruz and her team [192]. This data shows us that out of a total number of 874 grafts, 77.5% were used in glaucoma surgery, 5.2% in oculoplastic surgery, 5% in the treatment of defects of the cornea and sclera, 3.8% in reconstructions of the eyeball and 2.9% in covering orbital implants. The authors describe an increasing trend in the use of the sclera in reconstructions of eyelids, and conversely a decreasing trend in its use in the treatment of defects of the cornea and sclera.

CONCLUSION

As ensues from the studies presented above, over the course of the last approximately 80 years the sclera has been used in several different indications. Some of these are now practically obsolete (for example its use in retinal detachment surgery with an external approach), some rather rare (closure of corneal perforation), but in many other cases the sclera continues to be either the unequivocal method of choice (especially its use in glaucoma surgery, as well as treatment of retraction of the eyelids, extrusion of ocular implants or a scleral defect), or at least a relevant variant in the case of unavailability of otherwise frequently used material (for example the treatment of extensive defects of the eyelids or scleroplasty). Its main advantage is its good mechanical properties and low immunogenicity thanks to its high collagen content and only minimal content of blood vessels and cells. As a result, even despite the fact that it is now used relatively less frequently in comparison with other tissue banking products, it remains a useful material with further future potential for use in ophthalmic surgery.

REFERENCES

1. Costa LR. [Subconjunctival scleral grafts]. *Arch Soc Cir Hosp*. 1945 Dec;15:755-758. Spanish.
2. Larsson S. Treatment of perforated corneal ulcer by autoplasic scleral transplantation. *Br J Ophthalmol*. 1948 Jan;32(1):54-57.
3. Kurz J. Přeměna sklerální tkáně po implantaci do lidské rohovky. [Transformation of the scleral tissue following implantation in the human cornea]. *Cesk Oftalmol*. 1953 Oct;9(5):393-401. Czech.
4. Kok-Van Alphen CC. Corneal grafting of sclera, skin and mucous membrane. *Ophthalmologica*. 1951 Dec;122(6):382-384.
5. Paufique L, Moreau PG. [Scleromalacia perforans: histologic aspect, treatment by scleral transplant]. *Ann Ocul (Paris)*. 1953 Dec;186(12):1065-1076.
6. Kanagasundaram CR. Repair of perforating injury with a scleral graft. *Br J Ophthalmol*. 1959 Jul;43(7):440-441.
7. Obear MF, Winter FC. Technique of overlay scleral homograft. *Arch Ophthalmol*. 1964 Jun;71:837-838.
8. Payrau P, Remky H. [Scleroplasty with preserved sclera]. *Klin Monbl Augenheilkd Augenarztl Fortbild*. 1961 Jul;138:797-804.
9. Rodriguez-Vasquez F. New implant for retinal detachment surgery. *Am J Ophthalmol*. 1962;53:937-943.
10. Wilson FM. Homografts of preserved sclera in retinal detachment surgery. *Arch Ophthalmol*. 1964 Aug;72:212-218.
11. Knobloch WH, Cibis PA. Retinal detachment surgery with preserved human sclera. *Am J Ophthalmol*. 1965 Aug;60:191-204.
12. Stilma JS. Conjunctival excision or lamellar scleral autograft in 38 Moore's ulcers from Sierra Leone. *Br J Ophthalmol*. 1983 Jul;67(7):475-478.
13. Wang B, Wang Y, Ding J. [Repair of cornea-sclera defect by autogenous sclera graft from the same eye]. *Zhongguo Xiu Fu Chong Jian Wai Ke Za Zhi*. 1997 Jul;11(4):206-207.
14. Chaidaroon W, Sonthi W, Manassakorn A. Penetrating keratoplasty following scleral patch graft procedure. *J Med Assoc Thai*. 2004 Jan;87(1):53-58.
15. Prydal JI. Use of an autologous lamellar scleral graft to repair a corneal perforation. *Br J Ophthalmol*. 2006 Jul;90(7):924.
16. Levartovsky S, Springer A, Leiba H, Marcovich AL, Pollack A. Homologous scleral graft for corneal perforation in a child. *Cornea*. 2008 Feb;27(2):230-231.
17. Turner SJ, Johnson Z, Corbett M, Prydal J. Scleral autoplasty for the repair of corneal perforations: a case series. *Br J Ophthalmol*. 2010 May;94(5):669-670.
18. Sharma A, Mohan K, Sharma R, Nirankari VS. Scleral Patch Graft Augmented Cyanoacrylate Tissue Adhesive for Treatment of Moderate-Sized Noninfectious Corneal Perforations (3.5-4.5 mm). *Cornea*. 2013 Oct;32(10):1326-1330.

19. Prasher P. Use of an autologous lamellar scleral graft to repair a corneal perforation. *Int Ophthalmol*. 2014 Aug;34(4):957-960.
20. Jovanovic V, Jankov M, Nikolic L. Treatment of perforated cornea with an autologous lamellar scleral graft: histologic findings. *Arq Bras Oftalmol*. 2018 Jan-Feb;81(1):59-62.
21. Jyothi ELS, Krishna OG, Rao PL, Sekhar PR. Case series of scleral patch grafts during COVID-19 pandemic. *Indian J Ophthalmol*. 2021 Sep;69(9):2530-2532.
22. Karaca U, Usta G. The usability of lamellar scleral autograft in ocular perforation treatment. *Int Ophthalmol*. 2022 Feb;42(2):377-383.
23. Armstrong K, McGovern VJ. Scleromalacia perforans with repair grafting. *Trans Ophthalmol Soc Aust*. 1955;15:110-121.
24. Sevel D, Abramson A. Necrogranulomatous scleritis treated by an onlay scleral graft. *Br J Ophthalmol*. 1972 Nov;56(11):791-799.
25. de la Maza MS, Tauber J, Foster CS. Scleral grafting for necrotizing scleritis. *Ophthalmology*. 1989 Mar;96(3):306-310.
26. Saito N, Mori T, Kubota Y, Tochikubo T, Matsuhashi M, Komoto M. [A case of lamellar keratoplasty and autoscleroplasty combined with keratoepithelioplasty for Wegener's sclerocorneal ulcer]. *Nippon Ganka Gakkai Zasshi*. 1992 Aug;96(8):1061-1066. Japanese.
27. Kachmaryk M, Bouchard CS, Duffner LA. Bilateral fascia lata patch grafts in a patient with progressive scleromalacia perforans. *Ophthalmic Surg Lasers*. 1996 May;27(5):397-400.
28. Sangwan VS, Jain V, Gupta P. Structural and functional outcome of scleral patch graft. *Eye (Lond)*. 2007 Jul;21(7):930-935.
29. Aldasoro-Cáceres V, Aldasoro-Cáceres I, Pérez-Moreiras JV, Murrié-Fernández M, Ibáñez-Bosch R. Escleritis necrotizante c-ANCA positivo y escleritis múltiple compatible con Wegener ocular: tratamiento con rituximab [C-ANCA positive necrotising scleritis and multiple sclerosis compatible with ocular Wegener: treatment with rituximab]. *Arch Soc Esp Oftalmol*. 2014 Jan;89(1):31-34. Spanish.
30. Wen JC, Lam J, Banitt MR. Scleral Patch Graft With a Suture Reinforcement Technique in Surgical Management of Necrotizing Scleritis With Ectasia. *Cornea*. 2018 Jul;37(7):933-935.
31. Ryan SJ, Goldberg MF. Scleral surgery: selected indications and techniques. *Doc Ophthalmol*. 1971 May 14;29(2):261-287.
32. Gopal L, Badrinath SS. Autoscleral flap grafting: a technique of scleral repair. *Ophthalmic Surg*. 1995 Jan-Feb;26(1):44-48.
33. Oh JH, Kim JC. Repair of scleromalacia using preserved scleral graft with amniotic membrane transplantation. *Cornea*. 2003 May;22(4):288-293.
34. Ozcan AA, Bilgic E, Yagmur M, Ersöz TR. Surgical management of scleral defects. *Cornea*. 2005 Apr;24(3):308-311.
35. Esquenazi S. Autogenous lamellar scleral graft in the treatment of scleral melt after pterygium surgery. *Graefes Arch Clin Exp Ophthalmol*. 2007 Dec;245(12):1869-1871.
36. Jain V, Shome D, Natarajan S, Narverkar R. Surgically induced necrotizing scleritis after pterygium surgery with conjunctival autograft. *Cornea*. 2008 Jul;27(6):720-721.
37. de Farias CC, Sterlenich T, de Sousa LB, Vieira LA, Gomes JÁP. Randomized trial comparing multilayer amniotic membrane transplantation with scleral and corneal grafts for the treatment of scleral thinning after pterygium surgery associated with beta therapy. *Cornea*. 2014 Nov;33(11):1197-1204.
38. Polat N. Use of an Autologous Lamellar Scleral Graft to Repair a Scleral Melt After Mitomycin Application. *Ophthalmol Ther*. 2014 Dec;3(1-2):73-76.
39. Pujari A, Chaniyara MH, Sharma P, Sharma N. Necrotizing scleritis following uncomplicated strabismus surgery. *Indian J Ophthalmol*. 2020 Nov;68(11):2555-2557.
40. Rajagopal R, Matai HD, Gopal L, Susvar P, Bhende PS. Anterior staphyloma repair following trauma and surgery - A retrospective review of techniques and outcomes over 25 years. *Indian J Ophthalmol*. 2022 Aug;70(8):2967-2971.
41. Lamba PA, Shukla KN, Ganapathy M. Rhinosporidium granuloma of the conjunctiva with scleral ectasia. *Br J Ophthalmol*. 1970 Aug;54(8):565-568.
42. de Doncker RM, de Keizer RJ, Oosterhuis JA, Maes A. Scleral melting in a patient with conjunctival rhinosporidiosis. *Br J Ophthalmol*. 1990 Oct;74(10):635-637.
43. Lin CP, Wu YH, Chen MT, Huang WL. Repair of a giant scleral ulcer with a scleral graft and tissue glue. *Am J Ophthalmol*. 1991 Feb 15;111(2):251.
44. Lin CP, Tsai MC, Wu YH, Shih MH. Repair of a giant scleral ulcer with preserved sclera and tissue adhesive. *Ophthalmic Surg Lasers*. 1996 Dec;27(12):995-999.
45. Castelino AM, Rao SK, Biswas J, Gopal L, Madhavan HN, Kumar SK. Conjunctival rhinosporidiosis associated with scleral melting and staphyloma formation: diagnosis and management. *Cornea*. 2000 Jan;19(1):30-33.
46. Siatiri H, Mirzaee-Rad N, Aggarwal S, Kheirkhah A. Combined Tenonplasty and Scleral Graft for Refractory Pseudomonas Scleritis Following Pterygium Removal with Mitomycin C Application. *Ophthalmic Vis Res*. 2018 Apr-Jun;13(2):200-202.
47. Nakazawa M, Tamai M, Kiyosawa M, Watanabe Y. Homograft of preserved sclera for post-traumatic scleral staphyloma in Ehlers-Danlos syndrome. *Graefes Arch Clin Exp Ophthalmol*. 1986;224(3):247-250.
48. Barman M, Finger PT, Milman T. Scleral patch grafts in the management of uveal and ocular surface tumors. *Ophthalmology*. 2012 Dec;119(12):2631-2636.
49. Zahiti-Rashiti K, Meyer P, Goldblum D. Oral Mucosa as a Treatment Option in Perforating Scleral Necrosis after Proton Beam Radiation Therapy. *Klin Monbl Augenheilkd*. 2018 Apr;235(4):459-460.
50. Correa ZM, Huth B, Augsburger JJ. Scleral necrosis in patients with posterior uveal melanomas evaluated by transscleral fine needle aspiration biopsy and treated by 125I plaque. *Arq Bras Oftalmol*. 2018 Jul-Aug;81(4):330-335.
51. Rodríguez-Ares MT, Touriño R, Capeans C, Sánchez-Salorio M. Repair of scleral perforation with preserved scleral and amniotic membrane in Marfan's syndrome. *Ophthalmic Surg Lasers*. 1999 Jun;30(6):485-487.
52. Jain V, Natarajan S, Shome D, Gadgil D. Spectacle-induced ocular trauma: an unusual mechanism. *Cornea*. 2007 Jan;26(1):109-110.
53. Turaga K, Senthil S, Jalali S. Recurrent spontaneous scleral rupture in Marfan's syndrome. *BMJ Case Rep*. 2016 May 19.
54. Tarkkanen A, Vannas S. Corneoscleral grafting in the treatment of malignant melanoma of the limbus. *Ann Chir Gynaecol Fenn*. 1967;56(3):327-329.
55. Wilensky JT, Holland MG. A pigmented tumor of the ciliary body. *Arch Ophthalmol*. 1974 Sep;92(3):219-220.
56. Rummelt V, Naumann GO, Folberg R, Weingeist TA. Surgical management of melanocytoma of the ciliary body with extrascleral extension. *Am J Ophthalmol*. 1994 Feb 15;117(2):169-176.
57. Peyman GA, Apple DJ. Local excision of choroidal malignant melanoma. *Arch Ophthalmol*. 1974 Sep;92(3):216-218.
58. Shields JA, Eagle RC, Marr BP, Shields CL, Grossniklaus HE, Stulting RD. Invasive spindle cell carcinoma of the conjunctiva managed by full-thickness eye wall resection. *Cornea*. 2007 Sep;26(8):1014-1016.
59. Paufigue L, Charleux J. [Corneo-scleral graft for the treatment of dermoids of the limbus]. *Bull Soc Ophtalmol Fr*. 1967 May-Jun;67(5):463-467.
60. Burillon C, Durand L. Solid dermoids of the limbus and the cornea. *Ophthalmologica*. 1997;211(6):367-372.
61. Guo H, Huang K, Zhou D et al. [Acute leukemia child with ocular Kaposi's sarcoma after hematopoietic stem cell transplantation: a case report and literatures review]. *Zhonghua Xue Ye Xue Za Zhi*. 2013 May;34(5):445-448. Chinese.
62. Kobtan H. Use of autologous fascia lata as a natural biomaterial for tectonic support in surgically induced necrotizing scleritis. *Eye (Lond)*. 2015 Apr;29(4):580-584.
63. Yoo WS, Kim CR, Kim BJ et al. Successful Treatment of Infectious Scleritis by Pseudomonas aeruginosa with Autologous Perichondrium Graft of Conchal Cartilage. *Yonsei Med J*. 2015 Nov;56(6):1738-1741.
64. Kim JT, Kim KW, Mun S, Chun YS, Kim JC. Transplantation of autologous perichondrium with amniotic membrane for progressive scleral necrosis. *Ocul Surf*. 2019 Jul;17(3):571-577.
65. Alkharashi M, Dagi AF, Dagi LR. Pericardial patch graft repair of severe localized scleral thinning encountered during strabismus surgery. *J AAPOS*. 2017 Apr;21(2):156.
66. Bhalla M, Luk S, Jolly R, Ahmed F. Two-stage aqueous drainage tube implant with pericardial graft patch. *Eye (Lond)*. 2019 Sep;33(9):1509-1516.
67. Droustas K, Giachos I, Katsiampoula V, Kourti P, Kymionis G, Petrou P. Treatment of Progressive Scleromalacia Perforans by Tectonic Enhancement With Lyophilized Equine Pericardium. *Cornea*. 2021 May 1;40(5):648-651.
68. Zgolli HM, Mabrouk S, Fekih O, Malek I, Nacef L. Superficial Temporal Muscle Fascia Grafting: Successful Transplant of Surgical-induced Necrotizing Scleritis. *Middle East Afr J Ophthalmol*. 2021 Jan 19;27(4):238-240.

69. Lamarca-Mateu J, Salvador-Culla B, Gómez-Benlloch A, Barraquer RI. Oral mucosa for reconstructive surgery in a case of severe inflammatory necrotizing sclero-uveitis. *Indian J Ophthalmol.* 2020 Jan;68(1):221-223.
70. Finger PT, Reddy HS, Maheshwari A. Tarsconjunctival flap for scleromalacia related to treatments for conjunctival melanoma. *Am J Ophthalmol Case Rep.* 2023 Jan 18.
71. de Farias CC, Sterlenich T, de Sousa LD, Vieira LA, Gomes JÁP. Randomized trial comparing multilayer amniotic membrane transplantation with scleral and corneal grafts for the treatment of scleral thinning after pterygium surgery associated with beta therapy. *Cornea.* 2014 Nov;33(11):1197-1204.
72. Syed YA, Rapuano CJ. Umbilical amnion and amniotic membrane transplantation for infectious scleritis and scleral melt: A case series. *Am J Ophthalmol Case Rep.* 2021 Jan 23.
73. Naseripour M, Aghaei H, Sedaghat A, Kheirkhah A, Jaber R, Azma Z. Corneal Patch Graft: A New Approach for Scleral Necrosis Secondary to Plaque Radiotherapy. *Cornea.* 2016 Apr;35(4):565-568.
74. Burcu A, Yalniz-Akkaya Z, Özdemir ES, Özbek-Uzman S. Donor Cornea 74. Use in Scleral Surface Reconstruction. *Turk J Ophthalmol.* 2021 Aug 27;51(4):192-198.
75. Balestrazzi E, Mosca L, Blasi MA, Giannico MI, Balestrazzi A. Tectonic Keratoplasty to Restore the Bulbar Wall after Block Excision of Benign and Malignant Intraocular Tumors. *J Ophthalmol.* 2019 May 2.
76. Borley WE, Snyder AA. Surgical treatment of high myopia; the combined lamellar scleral resection with scleral reinforcement using donor eye. *Trans Am Acad Ophthalmol Otolaryngol.* 1958 Nov-Dec;62(6):791-801.
77. Miller WW, Borley WE. Surgical treatment of degenerative myopia. Scleral reinforcement. *Am J Ophthalmol.* 1964 May;57:796-804.
78. Thompson FB. A simplified scleral reinforcement technique. *Am J Ophthalmol.* 1978 Dec;86(6):782-790.
79. Ward B, Tarutta EP, Mayer MJ. The efficacy and safety of posterior pole buckles in the control of progressive high myopia. *Eye (Lond).* 2009 Dec;23(12):2169-2174.
80. Autrata R, Šenková K, Unčovská E, Řehůřek J. Skleroplastika při progresivní mopii u dětí: 20-leté výsledky. *Folia strabologica et neuroophthalmologica.* 2011;10(1):18-19. Czech.
81. Xue A, Zheng L, Tan G et al. Genipin-Crosslinked Donor Sclera for Posterior Scleral Contraction/Reinforcement to Fight Progressive Myopia. *Invest Ophthalmol Vis Sci.* 2018 Jul 2;59(8):3564-3573.
82. Gerinec A, Slezakova G. Posterior scleroplasty in children with severe myopia. *Bratisl Lek Listy.* 2001;102(2):73-78.
83. Chen M, Dai J, Chu R, Qian Y. The efficacy and safety of modified Snyder-Thompson posterior scleral reinforcement in extensive high myopia of Chinese children. *Graefes Arch Clin Exp Ophthalmol.* 2013 Nov;251(11):2633-2638.
84. Széll N, Boross A, Facskó A, Sohajda Z. Results with Posterior Scleral Reinforcement for Progressive Highly Myopic Children in Hungary. *Klin Monbl Augenheilkd.* 2022 Sep;239(9):1125-1131.
85. Miller HA. Scleral buckling with human preserved sclera in retinal detachment surgery. *Nippon Ganka Gakkai Zasshi.* 1966 Nov;70(11):2771-2778.
86. Pischel DK. The use of scleral transplants in the treatment of retinal detachment. *Trans Pac Coast Otoophthalmol Soc Annu Meet.* 1967;51:9-12.
87. Breffeilh LA. The use of human sclera as an encircling band for surgical treatment of retinal detachment. *South Med J.* 1971 Nov;64(11):1346-1347.
88. O'Gawa GM, Carey JD. Homologous scleral explant buckles in retinal detachment surgery. *Am J Ophthalmol.* 1974 Apr;77(4):505-508.
89. François J, Verbraeken H, Hanssens M. Scleral pockets and lyophilized sclera in retinal detachments. *Ophthalmologica.* 1979;179(3):153-157.
90. Poulsen CD, Peto T, Grauslund J, Green A. Epidemiologic characteristics of retinal detachment surgery at a specialized unit in Denmark. *Acta Ophthalmol.* 2016 Sep;94(6):548-555.
91. Roldán-Pallarés M, del Castillo Sanz JL, Susi SA, Refojo MF. Long-term complications of silicone and hydrogel explants in retinal reattachment surgery. *Arch Ophthalmol.* 1999 Feb;117(2):197-201.
92. Sheu SJ, Chou LC, Lee IY, Wang CC. Histopathology of polytetrafluoroethylene (Gore-tex) as a scleral buckle in humans. *Ophthalmic Surg Lasers Imaging.* 2001 May-Jun;32(3):245-247.
93. Shah PK, Karandikar SS, Narendran V. Scleral Autograft for Optic Nerve Head Pit Associated Chronic Maculopathy. *Ophthalmic Surg Lasers Imaging Retina.* 2017 Mar 1;48(3):251-254.
94. Helveston EM, Young LL. Human scleral patch for repair of exposed or extruded orbital implants. *Surg Forum.* 1968;19:490-492.
95. Helveston EM. Human bank scleral patch; for repair of exposed or extruded orbital implants. *Arch Ophthalmol.* 1969 Jul;82(1):83-86.
96. Helveston EM. A scleral patch for exposed implants. *Trans Am Acad Ophthalmol Otolaryngol.* 1970 Nov-Dec;74(6):1307-1310.
97. Zolli C, Shannon GM. Experience with donor sclera for extruding orbital implants. *Ophthalmic Surg.* 1977 Feb;8(1):63-70.
98. Soll DB. The use of sclera in surgical management of extruding implants. *Ophthalmology.* 1978 Aug;85(8):863-868.
99. McCarthy RW, Swann ES. Autogenous scleral graft in implant extrusion. *Ophthalmic Surg.* 1980 Oct;11(10):686-687.
100. Goldberg MF. A simplified scleral graft technique for covering an exposed orbital implant. *Ophthalmic Surg.* 1988 Mar;19(3):206-211.
101. Rubin PA. Enucleation, evisceration, and exenteration. *Curr Opin Ophthalmol.* 1993 Nov;4(5):39-48.
102. Kim J-H, Khwarg S-I, Choung HK, Yu YS. Management of porous polyethylene implant exposure in patients with retinoblastoma following enucleation. *Ophthalmic Surg Lasers Imaging.* 2004 Nov-Dec;35(6):446-452.
103. Axmann S, Paridaens D. Anterior surface breakdown and implant extrusion following secondary alloplastic orbital implantation surgery. *Acta Ophthalmol.* 2018 May;96(3):310-313.
104. Soll DB. Donor sclera in enucleation surgery. *Arch Ophthalmol.* 1974 Dec;92(6):494-495.
105. Oestreicher JH, Liu E, Berkowitz M. Complications of hydroxyapatite orbital implants. A review of 100 consecutive cases and a comparison of Dexon mesh (polyglycolic acid) with scleral wrapping. *Ophthalmology.* 1997 Feb;104(2):324-329.
106. Inkster CF, Ng SGJ, Leatherbarrow B. Primary banked scleral patch graft in the prevention of exposure of hydroxyapatite orbital implants. *Ophthalmology.* 2002 Feb;109(2):389-392.
107. Arat YO, Shetlar DJ, Boniuk M. Bovine pericardium versus homologous sclera as a wrapping for hydroxyapatite orbital implants. *Ophthalmic Plast Reconstr Surg.* 2003 May;19(3):189-193.
108. Yoon JS, Lew H, Kook KH, Lee SY. The use of autogenous sclera as wrapping material in hydroxyapatite implantation. *Ophthalmologica.* 2009;223(1):7-11.
109. Gawdat TI, Ahmed RA. Scleral shield: primary results of a new surgical technique in augmenting porous orbital implant protection. *Eur J Ophthalmol.* 2014 Nov-Dec;24(6):948-952.
110. Nadal J, Daien V, Jacques J, Hoa D, Mura F, Villain M. Evisceration with autogenous scleral graft and bioceramic implantation within the modified scleral shell: 133 cases over 17 years. *Orbit.* 2019 Feb;38(1):19-23.
111. Smith B, Lisman RD. Use of sclera and liquid collagen in the camouflage of superior sulcus deformities. *Ophthalmology.* 1983 Mar;90(3):230-235.
112. Gayre GS, Debacker C, Lipham W, Tawfik HA, Holck D, Dutton JJ. Bovine pericardium as a wrapping for orbital implants. *Ophthalmic Plast Reconstr Surg.* 2001 Sep;17(5):381-387.
113. Chow K, Satchi K, McNab AA. Combined use of bulbar conjunctival pedicle flap and labial mucous membrane graft for porous orbital implant exposure: Long-term outcome. *Orbit.* 2018 Aug;37(4):293-298.
114. Wang J-K, Liao S-L, Lai P-C, Lin LLK. Prevention of exposure of porous orbital implants following enucleation. *Am J Ophthalmol.* 2007 Jan;143(1):61-67.
115. Bodian M. Repair of ptosis using human sclera. *Am J Ophthalmol.* 1968 Mar;65(3):352-358.
116. Helveston EM, Wilson DL. A suture-reinforced scleral sling. Technique for suspension of the ptotic upper lid. *Arch Ophthalmol.* 1975 Aug;93(8):643-645.
117. Wasserman BN, Sprunger DT, Helveston EM. Comparison of materials used in frontalis suspension. *Arch Ophthalmol.* 2001 May;119(5):687-691.
118. Krásný J, Měřicka P. Možnosti řešení ptózy v dětství i v dospělosti [Treatment of ptosis in children and adults]. *Cesk Slov Oftalmol.* 1999 May;55(3):145-154. Czech.
119. Chen L, Li J, Zhang C, Li Y, Hou L, Ma J. Surgical Interventions for Congenital Ptosis: a Systematic Review and Meta-analysis of 14 Randomized Controlled Trials. *Aesthetic Plast Surg.* 2023 May 5.
120. Nakauchi K, Mito H, Mimura O. Frontal suspension for congenital ptosis using an expanded polytetrafluoroethylene (Gore-Tex®) sheet: one-year follow-up. *Clin Ophthalmol.* 2013;7:131-136.
121. Théra JP, Tiama JM, Konipo A, Napo A, Bamani S. Traitement du ptosis congénital dans un pays à faible revenu: suspension du

- releveur au muscle frontal par le polypropylène à l'Institut d'Ophthalmologie tropicale de l'Afrique (IOTA) [Treatment of congenital ptosis in a low-income country: polypropylene frontalis sling at the African Institute of Tropical Ophthalmology]. *J Fr Ophtalmol*. 2020 Feb;43(2):123-127. French.
122. Tenzel RR, Miller GR, Rubenzik R. Cicatricial upper lid entropion. Treated with banked scleral graft. *Arch Ophthalmol*. 1975 Oct;93(10):999-1000.
 123. Rubenzik R, Tenzel RR, Miller GR. Repair of cicatricial entropion of the upper eyelid. *Am J Ophthalmol*. 1975 Aug;80(2):302-303.
 124. Dryden RM, Soll DB. The use of scleral transplantation in cicatricial entropion and eyelid retraction. *Trans Sect Ophthalmol Am Acad Ophthalmol Otolaryngol*. 1977 Jul-Aug;83(4 Pt 1):669-678.
 125. Thommy CP. Scleral homograft inlay for correction of cicatricial entropion and trichiasis. *Br J Ophthalmol*. 1981 Mar;65(3):198-201.
 126. Zhang L, Hou Z, Li Y, Li D. Combined Surgical Strategies Adjusted to Clinical Evaluations for Severe Upper Eyelid Cicatricial Entropion and Trichiasis. *J Craniofac Surg*. 2023 Mar-Apr;34(2):764-767.
 127. Crawford JS, Easterbrook M. The use of bank sclera to correct lid retraction. *Can J Ophthalmol*. 1976 Oct;11(4):309-322.
 128. Cooper WC. The surgical management of the lid changes of Graves' disease. *Ophthalmology*. 1979 Dec;86(12):2071-2080.
 129. Flanagan JC. Retraction of the eyelids secondary to thyroid ophthalmopathy—its surgical correction with sclera and the fate of the graft. *Trans Am Ophthalmol Soc*. 1980;78:657-685.
 130. Doxanas MT, Dryden RM. The use of sclera in the treatment of dysthyroid eyelid retraction. *Ophthalmology*. 1981 Sep;88(9):887-894.
 131. Hurwitz JJ, Archer KF, Gruss JS. Treatment of severe lower eyelid retraction with scleral and free skin grafts and bipedicle orbicularis flap. *Ophthalmic Surg*. 1990 Mar;21(3):167-172.
 132. Mourits MP, Koornneef L. Lid lengthening by sclera interposition for eyelid retraction in Graves' ophthalmopathy. *Br J Ophthalmol*. 1991 Jun;75(6):344-347.
 133. Feldman KA, Putterman AM, Farber MD. Surgical treatment of thyroid-related lower eyelid retraction: a modified approach. *Ophthalmic Plast Reconstr Surg*. 1992;8(4):278-286.
 134. Olver JM, Rose GE, Khaw PT, Collin JR. Correction of lower eyelid retraction in thyroid eye disease: a randomised controlled trial of retractor tenotomy with adjuvant antimetabolite versus scleral graft. *Br J Ophthalmol*. 1998 Feb;82(2):174-180.
 135. Oestreicher JH, Pang NK, Liao W. Treatment of lower eyelid retraction by retractor release and posterior lamellar grafting: an analysis of 659 eyelids in 400 patients. *Ophthalmic Plast Reconstr Surg*. 2008 May-Jun;24(3):207-212.
 136. Park E, Lewis K, Alghoul MS. Comparison of Efficacy and Complications Among Various Spacer Grafts in the Treatment of Lower Eyelid Retraction: A Systematic Review. *Aesthet Surg J*. 2017 Jul 1;37(7):743-754.
 137. Wesley WE, McCord Jr CD. Transplantation of eye bank sclera in the Cutler-Beard method of upper eyelid reconstruction. *Ophthalmology*. 1980 Oct;87(10):1022-1028.
 138. Carroll RP. Entropion following the Cutler-Beard procedure. *Ophthalmology*. 1983 Sep;90(9):1052-1055.
 139. Berlin AJ, Carim M, Langston RH, Price RL. Scleral grafting in the management of lignant conjunctivitis. *Ophthalmic Surg*. 1982 Apr;13(4):288-291.
 140. Leone Jr CR. Periosteal flap for lower eyelid reconstruction. *Am J Ophthalmol*. 1992 Oct 15;114(4):513-514.
 141. Nesterov AP, Kolesnikova LN. Implantation of a scleral strip into the supraciliary space and cyclodialysis in glaucoma. *Acta Ophthalmol (Copenh)*. 1978 Oct;56(5):697-704.
 142. Freedman J. Scleral patch grafts with Molteno setons. *Ophthalmic Surg*. 1987 Jul;18(7):532-534.
 143. Lam SA, Cheuk W, Lai JS. Short-term results of using Lamellar Corneo-scleral patch graft for the Ahmed glaucoma valve implant surgery. *Yan Ke Xue Bao*. 1997 Sep;13(3):109-112.
 144. Aslanides IM, Spaeth GL, Schmidt CM, Lanzl IM, Gandham SB. Autologous patch graft in tube shunt surgery. *J Glaucoma*. 1999 Oct;8(5):306-309.
 145. Smith MF, Doyle JW, Ticeroy Jr JW. A comparison of glaucoma drainage implant tube coverage. *J Glaucoma*. 2002 Apr;11(2):143-147.
 146. Zalta AH. Long-term experience of patch graft failure after Ahmed Glaucoma Valve® surgery using donor dura and sclera allografts. *Ophthalmic Surg Lasers Imaging*. 2012 Sep-Oct;43(5):408-415.
 147. Choudhari NS, Neog A, Sharma A, Iyer GK, Srinivasan B. Our experience of fibrin sealant-assisted implantation of Ahmed glaucoma valve. *Indian J Ophthalmol*. 2013 Jan-Feb;61(1):23-27.
 148. Muir KW, Lim A, Stinnett S, Kuo A, Tseng H, Walsh MM. Risk factors for exposure of glaucoma drainage devices: a retrospective observational study. *BMJ Open*. 2014 May 2.
 149. Tsoukanas D, Xanthopoulos P, Charonis AC, Theodossiadi P, Kopsinis G, Filippopoulos T. Heterologous, Fresh, Human Donor Sclera as Patch Graft Material in Glaucoma Drainage Device Surgery. *J Glaucoma*. 2016 Jul;25(7):558-564.
 150. Wijsard MH, Haan M, Rietveld E, van Rijn LJ. Donor sclera versus bovine pericardium as patch graft material in glaucoma implant surgery and the impact of a drainage suture. *Acta Ophthalmol*. 2018 Nov;96(7):692-698.
 151. Pakravan M, Hatami M, Efsandiari H et al. Ahmed Glaucoma Valve Implantation: Graft-Free Short Tunnel Small Flap versus Scleral Patch Graft after 1-Year Follow-up: A Randomized Clinical Trial. *Ophthalmol Glaucoma*. 2018 Nov-Dec;1(3):206-212.
 152. Wang Y, Li X, Huang W et al. Partial thickness cornea tissue from small incision lenticule extraction: A novel patch graft in glaucoma drainage implant surgery. *Medicine (Baltimore)*. 2019 Mar.
 153. Kitnarong N, Srikulsasitorn B, Aurboonsong T. Glycerin-preserved Human-donor Corneoscleral Patch Grafts for Glaucoma Drainage Devices. *J Glaucoma*. 2020 Nov;29(11):1065-1069.
 154. Panda S, Khurana M, Vijaya L, George R, Balekudaru S. Comparison of conjunctiva-related complications between scleral and corneal patch grafts in Ahmed glaucoma valve implantation. *Indian J Ophthalmol*. 2023 Mar;71(3):881-7.
 155. Zeppa L, Romano MR, Capasso L, Tortori A, Majorana MA, Costagliola C. Sutureless human sclera donor patch graft for Ahmed glaucoma valve. *Eur J Ophthalmol*. 2010 May-Jun;20(3):546-51.
 156. Wigton E, Swanner JC, Joiner W et al. Outcomes of shunt tube coverage with glycerol preserved cornea versus pericardium. *J Glaucoma*. 2014 Apr-May;23(4):258-261.
 157. Lind JT, Shute TS, Sheybani A. Patch graft materials for glaucoma tube implants. *Curr Opin Ophthalmol*. 2017 Mar;28(2):194-198.
 158. Sheha H, Tello C, Al-Aswad LA, Sayed MS, Lee RK. Outcomes of the Shunt Tube Exposure Prevention Study: A Randomized Clinical Trial. *Ophthalmol Glaucoma*. 2019 Nov-Dec;2(6):392-401.
 159. Ahmed F, Mohamed NG. TAG (Tube and Graft) Sandwich Technique: A Novel Single-Stage Scleral Reinforcement and Aqueous Drainage Tube Implantation. *Case Rep Ophthalmol Med*. 2021 Jul 14.
 160. Singh M, Chew PTK, Tan D. Corneal patch graft repair of exposed glaucoma drainage implants. *Cornea*. 2008 Dec;27(10):1171-1173.
 161. Spierer O, Waisbourd M, Golan Y, Newman H, Rachmiel R. Partial thickness corneal tissue as a patch graft material for prevention of glaucoma drainage device exposure. *BMC Ophthalmol*. 2016 Feb 27.
 162. Chun YS, Kim KW, Kim JC. Autologous tragal perichondrium patch graft for ahmed glaucoma valve tube exposure. *J Glaucoma*. 2013 Dec.
 163. Stephens JD, Sarkisian SR Jr. The use of collagen matrix (Ologen) as a patch graft in glaucoma tube shunt surgery, a retrospective chart review. *F1000Res*. 2016 Aug 1.
 164. Gdih G, Jiang K. Graft-free Ahmed valve implantation through a 6 mm scleral tunnel. *Can J Ophthalmol*. 2017 Feb;52(1):85-91.
 165. Eslami Y, Azaripour E, Mohammadi M, Kiarudi MY, Fakhraie G, Zarei R et al. Single long scleral tunnel technique for prevention of Ahmed valve tube exposure. *Eur J Ophthalmol*. 2019 Jan;29(1):52-56.
 166. Melamed S, Ashkenazi I, Belcher DC 3rd, Blumenthal M. Donor scleral graft patching for persistent filtration bleb leak. *Ophthalmic Surg*. 1991 Mar;22(3):164-165.
 167. Clune MJ, Shin DH, Olivier MM, Kupin TH. Partial-thickness scleral-patch graft in revision of trabeculectomy. *Am J Ophthalmol*. 1993 Jun 15;115(6):818-820.
 168. Yan L, Zhang S, Dong Q. [The effect of heterogeneous scleral transplantation on hypotony after filtration surgery of glaucoma with mitomycin C]. *Zhonghua Yan Ke Za Zhi*. 1997 May;33(3):181-183.
 169. Kosmin AS, Wishart PK. A full-thickness scleral graft for the surgical management of a late filtration bleb leak. *Ophthalmic Surg Lasers*. 1997 Jun;28(6):461-468.
 170. Morris DA, Ramocki JM, Shin DH, Glover BK, Kim YY. Use of autologous Tenon's capsule and scleral patch grafts for repair of excessively draining fistulas with leaking filtering blebs. *Glaucoma*. 1998 Dec;7(6):417-419.
 171. Oyakhire JO, Moroi SE. Clinical and anatomical reversal of long-term hypotony maculopathy. *Am J Ophthalmol*. 2004 May;137(5):953-955.

172. Kompa S, Redbrake C, Arend O, Remky A. [Defect closure with scleral grafts]. *Klin Monbl Augenheilkd*. 2004 Oct;221(10):867-871.
173. Halkiadakis I, Lim P, Moroi SE. Surgical results of bleb revision with scleral patch graft for late-onset bleb complications. *Ophthalmic Surg Lasers Imaging*. 2005 Jan-Feb;36(1):14-23.
174. Harizman N, Ben-Cnaan R, Goldenfeld M, Levkovitch-Verbin H, Melamed S. Donor scleral patch for treating hypotony due to leaking and/or overfiltering blebs. *J Glaucoma*. 2005 Dec;14(6):492-496.
175. Au L, Wechsler D, Spencer F, Fenerty C. Outcome of bleb revision using scleral patch graft and conjunctival advancement. *J Glaucoma*. 2009 Apr-May;18(4):331-335.
176. King AJ, Rotchford AP. The use of a scleral micro-patch graft and fibrin glue to treat scleral flap defects following trabeculectomy. *Br J Ophthalmol*. 2009 Sep;93(9):1269-1270.
177. Mardelli PG, Mardelli ME, Bakkour Z. A Novel Hinged Scleral Patch Graft for the Repair of Overfiltration and Bleb Leaks. *J Glaucoma*. 2018 Apr;27(4):377-381.
178. Prokosch-Willing V, Lamparter J, Hassan SNU, Toshev AP, Pfeiffer N, Hoffmann EM. Results of an Adaptive Surgical Approach for Managing Late Onset Hypotony After Trabeculectomy With Mitomycin C. *J Glaucoma*. 2018 Apr;27(4):307-314.
179. O'Rourke M, Moran S, Collins N, Doyle A. Bleb reconstruction using donor scleral patch graft for late bleb leak and hypotony. *Eur J Ophthalmol*. 2021 May;31(3):1039-1046.
180. Rai P, Lauande-Pimentel R, Barton K. Amniotic membrane as an adjunct to donor sclera in the repair of exposed glaucoma drainage devices. *Am J Ophthalmol*. 2005 Dec;140(6):1148-1152.
181. Liu X, Law SK. Autologous Partial-thickness Scleral Flap and Donor Corneal Graft in Management of Tube Erosion of Glaucoma Drainage Device. *J Glaucoma*. 2019 Apr;28(4):347-351.
182. Zheng P, Pang X. Bleb-associated endophthalmitis treated by sclera patch graft, vitrectomy and endoscopic cyclophotocoagulation. *Chin Med J (Engl)*. 2012 Sep;125(18):3344-3345.
183. Thakur S, Ichhpujani P, Kumar S. Grafts in Glaucoma Surgery: A Review of the Literature. *Asia Pac J Ophthalmol (Phila)*. 2017 Sep-Oct;6(5):469-476.
184. Bochmann F, Kaufmann C, Kipfer A, Thiel MA. Corneal patch graft for the repair of late-onset hypotony or filtering bleb leak after trabeculectomy: a new surgical technique. *J Glaucoma*. 2014 Jan;23(1):e76-80.
185. Devloo S, Deghislage C, van Malderen L, Goethals M, Zeyen T. Non-penetrating deep sclerectomy without or with autologous scleral implant in open-angle glaucoma: medium-term results. *Graefes Arch Clin Exp Ophthalmol*. 2005 Dec;243(12):1206-1212.
186. Mousa ASG. Preliminary evaluation of nonpenetrating deep sclerectomy with autologous scleral implant in open-angle glaucoma. *Eye (Lond)*. 2007 Sep;21(9):1234-1238.
187. Naumann GO, Rummelt V. Block excision of cystic and diffuse epithelial ingrowth of the anterior chamber. Report on 32 consecutive patients. *Arch Ophthalmol*. 1992 Feb;110(2):223-227.
188. Forster RK. Corneoscleral block excision of postoperative anterior chamber cysts. *Trans Am Ophthalmol Soc*. 1995;93:83-97.
189. Rummelt V, Naumann GO. Blockexzision mit tektonischer Korneoskleralplastik wegen zystischer und/oder diffuser Epithelinvasion des vorderen Augenabschnitts - Bericht über 51 konsekutive Patienten (1980-1996) [Block excision with tectonic corneoscleroplasty for cystic and/or diffuse epithelial invasion of the anterior eye segment. Report of 51 consecutive patients (1980-1996)]. *Klin Monbl Augenheilkd*. 1997 Nov;211(5):312-323. German.
190. Thorisdottir RL, Sheikh R, Blohmé J, Stigmar G, Malmsjö M. A novel surgical technique employing donor sclera in strabismus surgery. *J AAPOS*. 2018 Jun;22(3):225-227.
191. Thorisdottir RL, Blohmé J, Malmsjö M. Clinical evidence supporting the use of donor sclera as spacer material in complicated cases of strabismus surgery - retrospective evaluation of surgical results in 117 patients with thyroid-associated ophthalmopathy or congenital strabismus. *Acta Ophthalmol*. 2019 Feb;97(1):74-79.
192. Sabater-Cruz N, Figueras-Roca M, Ventosa AG, Padró-Pitarch L, Tort J, Casaroli-Marano RP. Current clinical application of sclera and amniotic membrane for ocular tissue bio-replacement. *Cell Tissue Bank*. 2020 Dec;21(4):597-603.